

Appendix

GlossaryA-2
CreditsA-7

Glossary

A

acute triangle An acute triangle is a triangle with all three angles less than 90° .

altitude An altitude of a triangle is a line segment connecting a vertex to the line containing the opposite side and perpendicular to that side.

angle-angle-angle (AAA) similarity

The angle-angle-angle (AAA) similarity test says that if two triangles have corresponding angles that are congruent, then the triangles are similar. Because the sum of the angles in a triangle must be 180° , we really only need to know that two pairs of corresponding angles are congruent to know the triangles are similar.

angle bisector An angle bisector is a ray that cuts the angle exactly in half, making two equal angles.

C

central angle A central angle is an angle with its vertex at the center of a circle.

centroid The centroid of a triangle is the point where the three medians meet. This point is the center of mass for the triangle. If you cut a triangle out of a piece of paper and put your pencil point at the centroid, you could balance the triangle.

circle A circle is the set of all points in a plane that are equidistant from a given point in the plane, which is the center of the circle.

circumcenter The circumcenter of a triangle is the point where the three perpendicular bisectors meet. This point is the same distance from each of the three vertices of the triangles.

concave polygon A concave polygon is any polygon with an angle measuring more than 180° . Concave polygons look like they are collapsed or have one or more angles dented in.

concurrent When three or more lines meet at a single point, they are said to be concurrent. In a triangle, the three medians, three perpendicular bisectors, three angle bisectors, and three altitudes are each concurrent.

congruent Two figures are congruent if all corresponding lengths are the same, and if all corresponding angles have the same measure. Colloquially, we say they “are the same size and shape,” though they may have different orientation. (One might be rotated or flipped compared to the other.)

congruent triangles Congruent triangles are triangles that have the same size and shape. In particular, corresponding angles have the same measure, and corresponding sides have the same length.

converse Converse means the “if” and “then” parts of a sentence are switched. For example, “If two numbers are both even, then their sum is even” is a true statement. The converse would be “If the sum of two numbers is even, then the numbers are even,” which is not a true statement.

convex polygon A convex polygon is any polygon that is not concave.

coordinates Points are geometric objects that have only location. To describe their location, we use coordinates. We begin with a standard reference frame (typically the x- and y-axes). The coordinates of a point describe where it is located with respect to this reference frame. They are given in the form (x,y) where the x represents how far the point is from 0 along the x-axis, and the y represents how far it is from 0 along the y-axis. The form (x,y) is a standard convention that allows everyone to mean the same thing when they reference any point.

cosine If angle A is an acute angle in a right triangle, the cosine of A is the length of the side adjacent to angle A, divided by the length of the hypotenuse of the triangle. We often abbreviate this as $\cos A = (\text{adjacent})/(\text{hypotenuse})$.

cross section A cross section is the face you get when you make one slice through an object.

D

diameter A circle’s diameter is a segment that passes through the center and has its endpoints on the circle.

Glossary, cont'd.

E

edge An edge is a line segment where two faces intersect.

equilateral triangle An equilateral triangle is a triangle with three equal sides.

F

face A face is a polygon by which a solid object is bound. For example, a cube has six faces. Each face is a square.

frieze pattern A frieze pattern is an infinite strip containing a symmetric pattern.

G

glide reflection A glide reflection is a combination of two transformations: a reflection over a line followed by a translation in the same direction as the line.

H

hypotenuse The hypotenuse in a right triangle is the side of the triangle that is opposite to the right angle.

I

incenter The incenter of a triangle is the point where the three angle bisectors meet. This point is the same distance from each of the three sides of the triangle.

inscribed angle An inscribed angle is an angle whose vertex is on a circle and whose rays intersect the circle.

intercept An intercept is an intersection of a graph with one of the axes. An intersection with the horizontal axis is often referred to as an x-intercept, and an intersection with the vertical axis is often referred to as a y-intercept.

irregular polygon An irregular polygon is any polygon that is not regular.

isosceles trapezoid An isosceles trapezoid is a quadrilateral with one pair of parallel sides and congruent base angles, or it is a trapezoid with congruent base angles.

isosceles triangle An isosceles triangle is a triangle with two equal sides.

K

kite A kite is a quadrilateral that has two pairs of adjacent sides congruent (the same length).

L

line A line has only one dimension: length. It continues forever in two directions (so it has infinite length), but it has no width at all. A line connects two points via the shortest path, and then continues on in both directions.

line segment A line segment is the portion of a line lying strictly between two points. It has a finite length and no width.

line symmetry or **reflection symmetry** A polygon has line symmetry, or reflection symmetry, if you can fold it in half along a line so that the two halves match exactly. The folding line is called the line of symmetry.

M

median A median is a segment connecting any vertex of a triangle to the midpoint of the opposite side.

midline A midline is a segment connecting two consecutive midpoints of a triangle.

midline theorem The midline theorem states that a midline of a triangle creates a segment that is parallel to the base and half as long.

N

net A net is a two-dimensional representation of a three-dimensional object.

O

obtuse triangle An obtuse triangle is a triangle with one angle more than 90° .

orthocenter The orthocenter of a triangle is the point where the three altitudes meet, making them concurrent.

Glossary, cont'd.

P

parallel lines Parallel lines are two lines in the same plane that never intersect. Another way to think about parallel lines is that they are “everywhere equidistant.” No matter where you measure, the perpendicular distance between two parallel lines is constant.

parallelogram A parallelogram is a quadrilateral that has two pairs of opposite sides that are parallel.

perpendicular bisector The perpendicular bisector of a line segment is perpendicular to that segment and bisects it; that is, it goes through the midpoint of the segment, creating two equal segments.

plane A plane is a flat, two-dimensional object. We often represent a plane by a piece of paper, a blackboard, or the top of a desk. In fact, none of these is actually a plane, because a plane must continue infinitely in all directions and have no thickness at all. A plane can be defined by two intersecting lines or by three non-collinear points.

Platonic solid A Platonic solid is a solid such that all of its faces are congruent regular polygons and the same number of regular polygons meet at each vertex.

point A point specifies only location; it has no length, width, or depth. We usually represent a point with a dot on paper, but the dot we make has some dimension, while a true point has dimension 0.

polygon A polygon is a two-dimensional geometric figure with these characteristics:

It is made of straight line segments.

Each segment touches exactly two other segments, one at each of its endpoints.

It is closed—it divides the plane into two distinct regions, one inside and the other outside the polygon.

polyhedron A polyhedron is a closed three-dimensional figure. All of the faces are made up of polygons.

Pythagorean theorem The Pythagorean theorem states that if you have a right triangle, then the square built on the hypotenuse is equal to the sum of the squares built on the other two sides. $a^2 + b^2 = c^2$

Q

quadrilateral A quadrilateral is a polygon with exactly four sides.

R

radius The radius of a circle is the distance from the circle's center to a point on the circle, and is constant for a given circle.

ray A ray can be thought of as a half a line. It has a point on one end, and it extends infinitely in the other direction.

rectangle A rectangle is a quadrilateral with four right angles.

reflection Reflection is a rigid motion, meaning an object changes its position but not its size or shape. In a reflection, you create a mirror image of the object. There is a particular line that acts like the mirror. In reflection, the object changes its orientation (top and bottom, left and right). Depending on the location of the mirror line, the object may also change location.

regular polygon A regular polygon has sides that are all the same length and angles that are all the same size.

rhombus A rhombus is a quadrilateral that has all four sides congruent.

right triangle A right triangle is a triangle with one right (90°) angle.

rotation Rotation is a rigid motion, meaning an object changes its position but not its size or shape. In a rotation, an object is turned about a “center” point, through a particular angle. (Note that the “center” of rotation is not necessarily the “center” of the object or even a point on the object.) In a rotation, the object changes its orientation (top and bottom). Depending on the location of the center of rotation, the object may also change location.

rotation symmetry A figure has rotation symmetry if you can rotate (or turn) that figure around a center point by fewer than 360° and the figure appears unchanged.

Glossary, cont'd.

S

scalene triangle A scalene triangle is a triangle with all three sides unequal.

side-angle-side (SAS) congruence Side-angle-side (SAS) congruence states that if any two sides of a triangle are equal in length to two sides of another triangle and the angles between each pair of sides have the same measure, then the two triangles are congruent; that is, they have exactly the same shape and size.

side-angle-side (SAS) similarity The side-angle-side (SAS) similarity test says that if two triangles have two pairs of sides that are proportional and the included angles are congruent, then the triangles are similar.

side-side-side (SSS) congruence The side-side-side (SSS) congruence states that if the three sides of one triangle have the same lengths as the three sides of another triangle, then the two triangles are congruent.

side-side-side (SSS) similarity The side-side-side (SSS) similarity test says that if two triangles have all three pairs of sides in proportion, the triangles must be similar.

similar Two polygons are similar polygons if corresponding angles have the same measure and corresponding sides are in proportion.

similar triangles Similar triangles are triangles that have the same shape but possibly different size. In particular, corresponding angles are congruent, and corresponding sides are in proportion.

sine If angle A is an acute angle in a right triangle, the sine of A is the length of the side opposite to angle A divided by the length of the hypotenuse of the triangle. We often abbreviate this as $\sin A = (\text{opposite})/(\text{hypotenuse})$.

square A square is a regular quadrilateral.

symmetry A design has symmetry if you can move the entire design by either rotation, reflection, or translation, and the design appears unchanged.

T

tangent If angle A is an acute angle in a right triangle, the tangent of A is the length of the side opposite to angle A divided by the length of the side adjacent to angle A . We often abbreviate this as $\tan A = (\text{opposite})/(\text{adjacent})$.

tangram A tangram is a seven-piece puzzle made from a square. A typical tangram set contains two large isosceles right triangles, one medium isosceles right triangle, two small isosceles right triangles, a square, and a parallelogram.

theorem A theorem in mathematics is a proven fact. A theorem about right triangles must be true for every right triangle; there can be no exceptions. Just showing that an idea works in several cases is not enough to make an idea into a theorem.

translation Translation is a rigid motion, meaning an object changes its position but not its size or shape. In a translation, an object is moved in a given direction for a particular distance. A translation is therefore usually described by a vector, pointing in the direction of movement and with the appropriate length. In translation, the object changes its location, but not its orientation (top and bottom, left and right).

translation symmetry Translation symmetry can be found only on an infinite strip. For translation symmetry, you can slide the whole strip some distance, and the pattern will land back on itself.

transversal A transversal is a line that passes through (transverses) two other lines. We often consider what happens when the two other lines are parallel to each other.

trapezoid A trapezoid is a quadrilateral that has one pair of opposite sides that are parallel.

triangle inequality The triangle inequality says that for three lengths to make a triangle, the sum of the lengths of any two sides must be greater than the third length.

Glossary, cont'd.

V

van Hiele levels Van Hiele levels make up a theory of five levels of geometric thought developed by Dutch educators Pierre van Hiele and Dina van Hiele-Geldof. The levels are (0) visualization, (1) analysis, (2) informal deduction, (3) deduction, and (4) rigor. The theory is useful for thinking about what activities are appropriate for students, what activities prepare them to move to the next level, and for designing activities for students who may be at different levels.

vector A vector can be used to describe a translation. It is drawn as an arrow. The arrowhead points in the direction of the translation, and the length of the vector tells you the length of the translation.

Venn diagram A Venn diagram uses circles to represent relationships among sets of objects.

vertex A vertex is the point where two sides of a polygon meet.

Credits

Web Site Production Credits

Learning Math: Geometry is a production of WGBH Interactive and WGBH Educational Programming and Outreach for Annenberg/CPB.

Copyright 2002 WGBH Educational Foundation. All rights reserved.

Senior Producer

Ted Sicker

Curriculum Director

Denise Blumenthal

Content Developer

Michelle Manes, Mathematics Teacher and Education Consultant

Curriculum Developer

Anna Brooks

Project Manager

Sanda Zdjelar

Special Projects Assistant

Nina Farouk

Core Advisors

Suzanne Chapin

Srdjan Divac

Carol Findell

Bowen Kerins

Designers

Plum Crane

Lisa Rosenthal

Christian Wise

Web Developers

Joe Brandt

Kirsten Connelly

Bob Donahue

Peter Pinch

Online Video Segment Coordinator

Mary Susan Blout

Business Managers

Walter Gadecki

Joe Karaman

Unit Managers

Maria Constantinides

Adriana Sacchi

With the assistance of

Geoffrey Dixon

Stephani Roberts Lincoln

Yasmin Madan

Kate Smyres

Julie Wolf

Video Series Production Credits

Learning Math: Geometry is a production of WGBH Educational Foundation for Annenberg/CPB.

Executive Producer

Michele Korf

Senior Project Director

Amy Tonkonogy

Producer/Director

John Browne

Content Developer/Facilitator

Michelle Manes, Mathematics Teacher and Education Consultant

Advisors

Nicholas Branca, San Diego State University, CA

Miriam Leiva, University of North Carolina, Charlotte

Carol Malloy, University of North Carolina, Chapel Hill

Bill Masalski, University of Massachusetts, Amherst

Arthur Powell, Rutgers-Newark College of Arts and Sciences, NJ

Content Editor

Srdjan Divac, Harvard University, MA

On Location Consultants

Kenton G. Findell

Bowen Kerins

Editor

Miguel Picker

Additional Editing

Dickran H. Manoogian

Associate Producers

Irena Fayngold

Pamela Lipton

Credits, cont'd.

Project Manager

Sanda Zdjelar

Production Manager

Mary Ellen Gardiner

Post Production Associate Producer

Peter Villa

Location Coordinator

Mary Susan Blout

Teacher Liaison

Lisa Eure

Set Assistant

Elena Graceffa

Camera

Bill Charette
Lance Douglas
Larry LeCain
Steve McCarthy
Dillard Morrison
David Rabinovitz

Audio

Steve Bores
Chris Bresnahan
Charlie Collias
Jose Leon
Dennis McCarthy
Gilles Morin

Interns

Timothy Barney
Ravi Blatt
Joe Gudema
Justin Hartery
Aimee Jones

Design

Gaye Korbet
Daryl Myers
Alisa Placas

Content Graphics

Miguel Picker

On-line Editors

Mark Geffen
John O'Brien

Sound Mix

John Jenkins
Dan Lesiw

Original Music

Tom Martin

Additional Music

Concerto in G Major, J.B. de Boismortier

Soloist

John Tyson
Recorder

Frances Conover Fitch

Harpsichord

Jane Starkman, Kathryn Shaw

Violins

Jann Cosart

Viola

Alice Robbins

Violincello

Tom Coleman

Double Bass

Narrator

Jeff Loeb

Business Manager

Joe Karaman

Unit Manager

Maria Constantinides

Office Coordinators

Justin Brown
Laurie Wolf

Special Thanks:

Session 2. Triangles and Quadrilaterals: Boston's New Bridge

Kirk Elwell
Lead Field Engineer
Bechtel/Parsons Brinkerhoff

Session 3. Polygons: Three-Dimensional Printing

David Russell
Director of Engineering
Z Corporation

Marina Hatsopoulos
CEO
Z Corporation

Tom Clay
President
Z Corporation

Session 4. Parallel Lines and Circles: Three-Dimensional Maps

Seth Teller
Associate Professor
Lab for Computer Science, MIT

Hilary Karls
Student, MIT

Priya Agrawal
Student, MIT

Session 5. Dissections and Proof: Window Making and Proof

Jim Ialeggio
Architectural Detail In Wood

Session 6. The Pythagorean Theorem: The Pythagorean Theorem and 18th-Century Cranes

Rick Brown
Massachusetts College of Art

Wyly Brown

Session 7. Symmetry: Wooden Boats and Line Symmetry

Nat Benjamin
Gannon and Benjamin Marine
Railway

Session 8. Similarity: Similar Triangles and Radiation Therapy

Dr. Anita Mahajan
Radiation Oncologist
Tufts - New England Medical
Center

Maxon J. Buscher
Chief Dosimetrist
Tufts - New England Medical
Center

Jason K. Talkington
Medical Dosimetry Student
Tufts - New England Medical
Center

Sandra John-Baptiste, R.T.T.
Medical Dosimetry Student
Tufts - New England Medical
Center

Kevin O'Leary
Tufts - New England Medical
Center

Sean O'Leary
Tufts - New England Medical
Center

Session 9. Solids: Architecture and Solid Geometry

Ann Beha
Principal
Ann Beha Architects

Session 10: Classroom Case Studies

Rose Christiansen, Lincoln
School, Brookline, MA

Michelle Kurchian, Alice M.
Barrows Elementary School,
Reading, MA

Catalina Saenz, Roxbury
Preparatory Charter School,
Roxbury, MA

Heidi Weber, F.A. Day Middle
School, Newton, MA

Site Location

Ferryway School, Malden, MA